Bis-Michael Acceptors as Novel Probes to Study the Keap1/Nrf2/ARE Pathway

Ludovic J. Deny,^{†,§} Hussein Traboulsi,^{§,‡} André M. Cantin,^{§,‡} Éric Marsault,^{§,||} Martin V. Richter,^{§,‡} and Guillaume Bélanger^{*,†,§}

[†]Département de Chimie, Université de Sherbrooke, 2500 Boulevard Université, Sherbrooke, Québec J1K 2R1, Canada

[§]Institut de Pharmacologie de Sherbrooke, Université de Sherbrooke, Sherbrooke, Québec J1H 5N4, Canada

[‡]Département de Médecine, Service de Pneumologie et Centre de Recherche Clinique, Centre Hospitalier de l'Université de Sherbrooke, Sherbrooke, Québec J1H 5N4, Canada

^{||}Département de Pharmacologie, Université de Sherbrooke, Sherbrooke, Québec J1K 2R1, Canada

S Supporting Information

ABSTRACT: Nuclear factor erythroid 2-related factor 2 (Nrf2) is a master regulator that promotes the transcription of cytoprotective genes in response to oxidative/electrophilic stress. Various Michael-type compounds were designed and synthesized, and their potency to activate the Keap1/Nrf2/ARE pathway was evaluated. Compounds bearing two Michael-type acceptors proved to be the most active. Tether length and rigidity between the acceptors was crucial. This study will help to understand how this feature disrupts the interaction between Keap1 and Nrf2.

INTRODUCTION

Oxidative/electrophilic stress (OES) is the consequence of an abnormal production of reactive oxygen and nitrogen species and/or electrophiles in the cells.¹ OES stress is linked to over 200 diseases and conditions such as cancers, diabetes, pulmonary infections, and neurodegenerative disorders. The Keap1/Nrf2/ARE pathway plays a central role in the cellular response to OES because the antioxidant response element (ARE) is an essential component of the cellular antioxidant defense and is under the transcriptional control of nuclear factor erythroid 2-related factor 2 (Nrf2).^{2–4}

Because Kelch like ECH associated protein 1 (Keap1) regulates Nrf2-mediated transcription, the former is recognized as a therapeutic target with high potential. Indeed, the genes targeted by the Keap1/Nrf2/ARE pathway are involved in the synthesis and conjugation of glutathione and antioxidant proteins and enzymes.⁵ Understanding the Keap1/Nrf2/ARE pathway has thus become extremely important and the focus of drug discovery programs recently initiated by several research groups.^{6–18}

In the basal state, Nrf2 is sequestered in the cytosol via its interaction with a Keap1 dimer. When this Keap1–Nrf2 complex binds to the Keap1-dependent E3-ubiquitin ligase (Cullin 3), Nrf2 is then degraded by the proteasome and the Keap1 dimer is free to bind to a newly translated Nrf2. This maintains low concentrations of free Nrf2 in the nucleus and

transcription of genes related to ARE at basal levels. When Keap1 is targeted by small molecules, the conformation of the Keap1 dimer is modified and its interactions with Nrf2 are tighter. Accordingly, the ubiquitination of Nrf2 is stopped and newly synthesized Nrf2 can accumulate in the nucleus and turn on the expression of cytoprotective genes. According to Dinkova-Kostova et al., this mode of action is proper to electrophilic inducers.¹⁹

Keap1 comprises five domains: (1) the N-terminal region, (2) the broad complex, tramtrack, bric-a-brac (BTB) domain (exposing Cys151, responsible for the formation of the Keap1 dimer), (3) the intervening region (IVR, exposing Cys273, Cys288, and six others), (4) the Kelch repeat domain (responsible for the recognition of Nrf2), and (5) the C-terminal region.²⁰ Among strategies used to disrupt the Keap1–Nrf2 interaction, noncovalent protein–protein interaction inhibitors have recently been reported.^{6–10} Several crystal structures reveal that these molecules target the binding site of Nrf2 on Keap1 (Kelch repeat domain).^{7–9} Another strategy consists of using electrophiles from synthetic or natural sources.^{11–18} The latter proved particularly successful (Figure 1) and are believed to target some of the cysteine (Cys) residues of Keap1. Among them, Cys151 (BTB domain) as well

Received: July 28, 2016

Published: September 28, 2016

2). For β -ketoesters bearing an unsaturation (cf. compounds **10** and **12**), final oxidation gave best results using DDQ.

Scheme 2^a

5

a
 72%

6

b, c, d
 or b, e

R

7 R = H (57%)
 8 R = C₆H₁₃ (34%)
 9 R = Me (39%)
 10 R = allyl (39%)
 11 R = Bn (40%)
 12 R = propargyl (34%)
 13 R = CH₂CF₃ (7%)
 14 R = CH₂CO₂tBu (19%)
 15 R = CH₂CN (9%)
 16 R = CH₂C(O)Me (4%)
 17 R = (CH₂)₇CO₂Me (16%)

■ SYNTHESIS

To study the influence of the ester portion, transesterification²⁹ of **6** using allyl alcohol or *tert*-butanol as the solvent gave mixtures of esters. After oxidation, **7** and **18** or **7** and **19**, respectively, could be isolated (Scheme 3). We also prepared the corresponding acid **20** by hydrolysis of unsaturated keto-ester **7**.¹³

To test the influence of the electrophilicity of the Michael acceptor, several variations were made (Scheme 4). First, a reduction of **7** in Luche conditions afforded the alcohol **21**.³⁰ Chloro- and iodoenones **23** and **24** were prepared using known procedures.^{31,32} Morita–Baylis–Hillman reaction on **22** with paraformaldehyde gave the allylic alcohol **25**.³³ Further transesterification afforded **26**.³⁴ Classic methoxycarbonylation of enone **22** provided keto-ester **27**, which, upon α -methylation, afforded **28** as a racemic mixture.³⁵ On the

Scheme 3^a

^aReagents and conditions: (a) DMAP, 4 Å MS, ROH, reflux; (b) PhSeBr, pyridine, CH₂Cl₂, 0 °C; (c) H₂O₂, CH₂Cl₂, 0 °C; (d) K₂CO₃, DDQ, 1,4-dioxane, rt; (e) KOH, H₂O/MeOH, reflux.

Scheme 4^a

^aReagents and conditions: (a) CeCl₃, NaBH₄, MeOH, rt; (b) PhI(OAc)₂, pyridine-HCl, CH₂Cl₂, rt; (c) I₂, pyridine, CCl₄, rt; (d) paraformaldehyde, NaHCO₃, imidazole, THF, rt; (e) 6, Et₃N, toluene, Dean–Stark; (f) NaH, Me₂CO₃, 1,4-dioxane, reflux; (g) MeI, K₂CO₃, acetone, 40 °C; (h) SeO₂, AcOH, *t*-BuOH, reflux.

other hand, oxidation of keto-ester 27 by selenium dioxide gave dienone 29.³⁶

Bis-Electrophilic Compounds. To attach two Michael acceptors on the same molecule and assess the influence of rigidity on activity, we used tethers of different lengths (1–4 carbons, 31,32) and rigidity (alkane, alkene 33, and alkyne 34, Scheme 5). These tethers were incorporated by bis-alkylation

Scheme 5^a

^aReagents and conditions: (a) LDA, R-X, rt; (b) PhSeBr, pyridine, CH₂Cl₂, 0 °C; (c) H₂O₂, CH₂Cl₂, 0 °C; (d) DDQ, 1,4-dioxane, rt; (e) DDQ, K₂CO₃, 1,4-dioxane, rt.

with the dianion of 6. No significant amount of single alkylation biproducts was observed. The ensuing tandem oxidations afforded the desired products 30–34, albeit in low yield. A series of compounds bearing a dioxolane instead of the *gem*-dimethyl was also prepared for comparison (35, 36). Finally, unsymmetrical bis-electrophilic compound 37 was obtained by oxidation of 26 with DDQ.

RESULTS AND DISCUSSION

ARE Inducing Activity and Cytotoxicity. To evaluate the ARE-inducing activity, all compounds, at concentrations ranging from 0 to 100 μM, were incubated with ARE-Luc-transfected human embryonic kidney cells (HEK-293T) for 24 h. Suforaphane (10 μM) was used as a positive control. Also, NAD(P)H:quinone oxidoreductase (NQO1) induction was measured by quantitative polymerase chain reaction (qPCR) in RAW 264.7 cells using sulforaphane as a positive control. In addition, cytotoxicity was evaluated for each molecule in a lactate dehydrogenase (LDH) assay in HEK-293T cells. The data are reported as mean ± SEM based on three separated experiments.

Lipophilicity As an Off-Target Indicator. Compounds lacking any substitution at the C4 position (see 7, with two hydrogens instead of *gem*-dimethyl) were too electrophilic.

Difficulties in their isolation and purification compromised the validity of their evaluation.

To counterbalance for the level of reactivity of these electrophiles, steric hindrance around C4 was increased. We chose commercially available ketone **1** as a starting point for comparison. Analogues with a dioxolane moiety (**3**, **4**, **35**, and **36**) showed moderate activity and significant toxicity at the higher concentrations (Figure 2).

Figure 2. (A) Fold induction of ARE-luciferase gene expression; (B) % cell mortality induced by compounds **3**, **4**, **35**, **36**, and **7** at 10, 20, 50, and 100 μ M, respectively.

We hypothesized that toxicity was due to the hydrophilicity associated with the dioxolane moiety.^{14,37} To test this hypothesis, we increased the compounds' lipophilicity. Replacing the dioxolane with a *gem*-dimethyl provided an increased activity with no toxicity even at high concentrations (e.g., **7**). Further exploration of the substitution at C6 was essential to generate unambiguous biological data (Table 1). The single electrophiles showed better activities at 100 μ M (compounds **7**–**12**), with no apparent toxicity. They were much more lipophilic than their dioxolane analogues according to their clogP values, which tends to support our hypothesis.

Importance of Electrophilicity. To induce the transcription of genes controlled by ARE, a first cysteine nucleophilic addition to the electrophilic compounds has to occur. Analogues **7**–**12** suggest that unsaturated keto esters are indeed suitable electrophiles to study Keap1 reactivity. Replacement of the methyl ester by larger groups (allyl **18** or *t*-Bu **19**) gave a slight increase in activity, although accompanied by increased toxicity, whereas the corresponding acid (**20**) retained most of the activity. Replacement of the methyl ester by a chloride (**23**) had no significant effect, but the

Table 1. Fold-Induction of ARE-Luciferase Gene Expression^a

Compound	1 μ M	Fold-induction at: ^b		clogP ^c
		20 μ M	100 μ M	
3	1.40 $\pm$ 0.25	2.19 $\pm$ 0.19	20.19 $\pm$ 1.10	2.49
4	0.98 $\pm$ 0.04	2.49 $\pm$ 0.08	N/A	-0.16
7	1.36 $\pm$ 0.34	6.46 $\pm$ 4.26	50.58 $\pm$ 12.39	1.82
8	1.15 $\pm$ 0.04	2.00 $\pm$ 0.43	64.65 $\pm$ 14.31	4.98
9	1.17 $\pm$ 0.11	2.75 $\pm$ 0.22	4.60 $\pm$ 0.10	2.34
10	1.34 $\pm$ 0.51	2.05 $\pm$ 0.06	25.32 $\pm$ 4.03	2.91
11	0.95 $\pm$ 0.12	0.92 $\pm$ 0.04	48.53 $\pm$ 14.74	3.91
12	1.14 $\pm$ 0.12	1.48 $\pm$ 0.44	34.16 $\pm$ 2.62 (7%)	2.39
13	0.54 $\pm$ 0.07	1.54 $\pm$ 0.22	17.90 $\pm$ 6.18 (11%)	2.58
14	0.65 $\pm$ 0.09	2.00 $\pm$ 0.17	1.75 $\pm$ 0.18	3.26
15	1.03 $\pm$ 0.39	2.22 $\pm$ 1.13	17.13 $\pm$ 2.52	1.40
16	0.49 $\pm$ 0.06	2.20 $\pm$ 0.07	8.71 $\pm$ 1.01	1.72
17	0.68 $\pm$ 0.08	1.24 $\pm$ 0.09	5.96 $\pm$ 1.17	2.30
18	1.39 $\pm$ 0.27	4.36 $\pm$ 0.50	94.79 $\pm$ 22.67 (17%)	2.59
19	1.05 $\pm$ 0.05	3.89 $\pm$ 0.39	106.40 $\pm$ 20.32 (16%)	3.06
20	2.00 $\pm$ 0.19	3.11 $\pm$ 0.73	29.56 $\pm$ 1.28	1.39
23	1.33 $\pm$ 0.18	4.42 $\pm$ 0.69	43.41 $\pm$ 3.32	2.34
24	1.22 $\pm$ 0.16	1.16 $\pm$ 0.23	17.35 $\pm$ 0.47	2.71
25	0.72 $\pm$ 0.05	0.77 $\pm$ 0.17	1.46 $\pm$ 0.13	0.87
26	2.98 $\pm$ 0.36	3.79 $\pm$ 1.10	4.02 $\pm$ 0.18	3.40
21	1.00 $\pm$ 0.07	1.85 $\pm$ 0.37	1.29 $\pm$ 0.34	1.91
27	1.05 $\pm$ 0.11	3.20 $\pm$ 0.35	41.93 $\pm$ 7.94	1.43
28	1.47 $\pm$ 0.13	2.43 $\pm$ 0.09	4.56 $\pm$ 0.41	1.95
29	1.35 $\pm$ 0.21	3.39 $\pm$ 0.41	10.32 $\pm$ 0.72	1.96
30	4.49 $\pm$ 0.43	3.60 $\pm$ 0.48	101.10 $\pm$ 15.76 (24%)	3.79
31	4.71 $\pm$ 1.11	247.70 $\pm$ 17.80	Not tested	4.51
32	8.23 $\pm$ 0.38	Not tested	Not tested	5.04
33	15.20 $\pm$ 0.79	Not tested	Not tested	4.56
34	52.46 $\pm$ 3.01	Not tested	Not tested	4.38
35	0.94 $\pm$ 0.03	2.76 $\pm$ 0.15	15.94 $\pm$ 0.67 (12 %)	-1.20
36	1.00 $\pm$ 0.04	1.05 $\pm$ 0.15	30.91 $\pm$ 2.14 (9%)	-0.48
37	2.02 $\pm$ 0.40	3.78 $\pm$ 0.56	121.1 $\pm$ 20.60 (15%)	3.28

^aSulforaphane has a fold-induction of 9.80 $\pm$ 1.00 at 10 μ M. ^b% Mortality evaluated in the LDH assay are reported when over 5%. ^cPredicted values by ChemDraw 12.0.

iodide derivative (**24**) was not reactive enough. This effect was also observed with other poor electrophiles, such as those lacking the ketone (**21**) or the ester (**25**, **26**, and **28**), with the exception of compound **27**. This comes as a surprise because it is hard to evaluate the combined participation of the carbonyls in **27** while we expected **29** to be much more potent.³⁸

Boost in Activity with a Second Michael Acceptor. To take advantage of the numerous cysteine residues of Keap1,³⁹ we tested compounds with two Michael-type acceptors. Right

away, this second class of molecules turned out to be very potent (30, 31, and 38), with an astonishing 167-fold increase from 7 to 31 at 10 μ M (Figure 3). Even though solubility and

Figure 3. Docking of 31 in BTB domain of Keap1 using PyMol and NRGsuite.

toxicity became an issue at higher concentrations, the study of similar compounds provided new insights about the target. First, the addition of a second unsaturated keto-ester enhanced the fold-induction drastically. Second, the spatial position between the two electrophilic parts is important to modulate activity. Indeed, going from a simple methylene (30) to a propane linker (31) resulted in a ca. 70-fold increase (at 20 μ M), whereas an additional factor 2 was gained with a butane tether (32).

So far, it was not clear whether the tether length optimization reflected an optimal distance between the two Michael acceptors for reaction with two cysteine residues in the binding site or the result of a gain in hydrophobic interactions. To answer this, we decided to rigidify the tether. From butane (32) to *E*-butene (33) to butyne (34) tethers, increased rigidity resulted in a significant increase in activity, with compound 34 being the best compound so far (50-fold-induction at only 1 μ M). These results strongly suggest an optimal distance and conformation between the Michael acceptors for interaction with Keap1.

These results raise the question whether the second Michael acceptor on 30–34 acts as a cysteine trap. It turns out that the Michael acceptors on these molecules are similar to the α,β' -unsaturated β -cyanoketone of CDDO, which was co-crystallized in the BTB domain of Keap1.²⁰ We have thus docked our molecules in the crystal structure and set the distance between the nucleophilic sulfur and the electrophilic carbon to 1.8 Å as in CDDO. Interestingly, there is no other cysteine residue in the active site of the BTB domain that could act as a partner for a second nucleophilic addition. In return, the ketone and the ester functions were able to make hydrogen bonds with other residues in the active site and within the electronic density area. Hence, in the absence of a crystal of Keap1 with any of compounds 30–34, we cannot answer the question on the nature of the interaction of the second Michael acceptor with Keap1. As suggested by docking, it could be a H-bond in the BTB domain but it could also be that these molecules rather target the IVR region. Unfortunately, this region of Keap1 has not been crystallized yet and we cannot answer this hypothesis clearly at this point.

Added Functionality Reduces Activity. To prove whether only an H-bond acceptor is needed in addition to

the electrophile, we installed different chemical functions (13–17). None of these compounds showed activity. Even 17 is particularly inactive, although it presents a distance between the ester side chain and the Michael-type acceptor that resembles the distance between Michael-type acceptors in 30. Having H-bond acceptors on the side chain (14–17) is even more detrimental than simple alkyl side chains (8, 10–12), suggesting that this part of the molecule also plays a role in lipophilic interactions with the active site.

NQO1 Expression. Michael-type acceptors are listed as pan assay interference compounds (PAINs).⁴⁰ To address this eventuality in the case of the present class of molecules, we measured the expression of NQO1, a Nrf2 positively regulated gene, by qPCR in RAW 264.7 cells for sulforaphane and 14 compounds of interest at 10 μ M (Table S1, Supporting Information).^{5,11,13,15,18} We observed the same tendencies as those observed in the luciferase ARE reported assay (Figure 4).

Figure 4. Fold induction of NQO1 expression by sulforaphane (SFN), compounds 8, 17, 27, 32, 33, and 34 at 2, 5, and 10 μ M.

Indeed, an aliphatic side chain (8) is preferred over a functionalized one (17), the addition of a Michael acceptor enhances activity (cf. 8 and 32), and an increased rigidity between Michael acceptors is beneficial (cf. 32, 33, and 34). This confirms that the observed activity is indeed dependent on the Nrf2 pathway and not on a promiscuous activity. Moreover, these Michael acceptors are much more potent than the naturally occurring sulforaphane.

CONCLUSION

In this study, we report herein the synthesis and biological evaluations of 32 Michael acceptors toward the Keap1/Nrf2/ARE pathway for the transcription of antioxidant cellular responses. This study gives several insights on how reversible covalent modulators interact with Keap1. The electrophilic part of the molecule has to be sufficiently lipophilic to show some activity. Furthermore, the electrophile has to be powerful enough to react with the targeted cysteine, an essential element in the design of Keap1 ligands to stop the ubiquitination and subsequent degradation of Nrf2. The 1-methoxycarbonyl-3,3-dimethylcyclohex-2-enone motif answered these two criteria. Addition of a second Michael acceptor had a crucial impact on activity, up to a 50-fold-induction at only 1 μ M. The distance between the Michael acceptors as well as the rigidity of the tether were optimized to generate our best compound (34) in this family.

Our results suggest that the role of the second Michael acceptor is unlikely to be a hydrogen bond acceptor as none of the compounds with heteroatoms at the end of the side chain showed activity. We could hypothesize that the second Michael acceptor rather serves as a second cysteine trap. This remains to be proven, but the consequences of such a hypothesis, should it

turn out to be true, are of high importance. Indeed, trapping of several cysteine residues would imply that the BTB domain is not the target but probably the IVR region of Keap1.

We believe our study will serve to increase the understanding of this target toward the generation of new drug leads. Our data also highlight the importance to design appropriate electrophiles toward covalent drug. Investigations on the nature and exact role of the second electrophile are in progress, and the results will be published in due course.

EXPERIMENTAL SECTION

Biology: Cell lines. Human embryonic kidney 293 cells (HEK-293T; ATCC CRL-1573) and the murine macrophage cell line (RAW 264.7; ATCC TIB-71) were obtained from American Type Culture Collection (Manassas, VA). Cells were cultured using standard methods at 37 °C in 5% CO₂–95% air in Dulbecco's Modified Eagle Medium (DMEM) (Wisent, Canada) supplemented with 10% fetal bovine serum (FBS), antibiotics (streptomycin at 100 mg/mL and penicillin at 100 U/mL), 5% sodium pyruvate, 5% nonessential amino acids, and 2 mM L-glutamine (complete DMEM).

Luciferase Assays. The day before transfection, human embryonic kidney HEK 293T cells (1×10^5) were plated in 24-well tissue culture-treated plates. Cells were transiently transfected with 0.1 μ g of ARE-luc (Nrf2 response element) plasmid using FuGENE6 transfection reagent (Promega, Madison, WI; E2691). The day after transfection, cells were treated with Keap1 ligands at different concentrations or with sulforaphane at 10 μ M. After 24 h, cell lysates were then assayed for luciferase activity using the luciferase assay system from Promega (E1500).

Real-Time PCR Analyses. RAW 264.7 cells were treated with 2, 5, and 10 μ M of different compounds or with 10 μ M of sulforaphane for 5 h. Total RNA was then extracted using RiboZol reagent according to the manufacturer's protocol. A total of 500 ng of the resulting RNA was then reverse transcribed using iScript reverse transcription supermix for the reverse transcriptase quantitative PCR (RTqPCR) kit (Bio-Rad). Real-time PCRs were performed with the iQ SYBR green supermix (Bio-Rad). Amplification plots were generated using the Rotorgene 6000 Application software version 1.7 (Corbett Research), and fold induction was calculated using the threshold cycle (2 $^{-\Delta\Delta C_T}$) method and using 18S expression for normalization. Primer sequences are

Human/mouse 18S:

FW: 5'-AGGAATTGACGGAAGGGCAC-3'

RV: 5'-GGACATCTAAGGGCATCACA-3'

Mouse NQO1:

FW: 5'-TCACAGGTGAGCTGAAGGAC-3'

RV: 5'-CTTCCAGCTTCTTGTTGTCG-3'

Cell Viability Assay. The cytotoxicity of Keap1 ligands on cells was determined using the LDH assay. Briefly, HEK 293T cells were treated with increasing concentrations of Keap1 ligands. After 24 h, supernatants were collected and cells were lysed in Triton 1%. To measure lactate dehydrogenase (LDH) activity, buffer containing L-(+)-lactic acid and nicotinamide adenine dinucleotide (NAD) (Sigma-Aldrich; L-1750 and N7004, respectively) was added to the supernatant and the cells lysate, followed by measurement of absorbance at 340 nm.

Chemistry: General. All reactions requiring anhydrous conditions were conducted in flame-dried glassware under a dry nitrogen or argon atmosphere. THF was distilled from Na and benzophenone under nitrogen immediately prior to use. 1,4-Dioxane was distilled from Na and benzophenone under argon and kept on 4 Å molecular sieves. DCM, MeOH, toluene, *i*-Pr₂NH, Et₃N, and pyridine were distilled from CaH₂ under nitrogen at atmospheric pressure immediately prior to use. All other required fine chemicals were used directly without purification. All reference to "water" correspond to deionized water. All references to "brine" refer to a saturated aqueous sodium chloride solution. Melting points were determined on a Thomas-Hoover capillary melting point apparatus and are uncorrected. All new

compounds showed chemical purity of $\geq 95\%$ as assessed by ¹H NMR. All NMR spectra were recorded on an AV300 Bruker (300 MHz for ¹H and 75 MHz for ¹³C) or an AS400 variant (400 MHz for ¹H and 100 MHz for ¹³C). Chemical shifts are referenced to δ 7.26 signal of CHCl₃ (¹H NMR) and 77.16 signal of CDCl₃ (¹³C NMR) as internal standards for deuterated chloroform and to δ 5.30 signal of CH₂Cl₂ (¹H NMR) and δ 53.52 signal of CD₂Cl₂ (¹³C NMR) as internal standards for deuterated DCM. Data for proton spectra are reported as follows: chemical shift in ppm (multiplicity [singlet (s), doublet (d), triplet (t), quartet (q), and multiplet (m)]), integration, coupling constants [Hz]. Carbon spectra were recorded with complete proton decoupling, and the chemical shifts are reported in ppm. High-resolution mass spectrometry data were obtained by the ESI-Q-ToF (maXis). TLC was conducted with precoated 60 Å 250 μ m silica gel plates with F-254 indicator and visualized using a combination of UV and potassium permanganate staining. Flash column chromatography was performed using silica gel (230–400 mesh). IR spectra were recorded with a FTIR instrument by applying substrates neat.

Usual Reaction Workup and Purification. After addition of the indicated aqueous solution, layers were separated. The aqueous phase was extracted with the indicated solvent, and the combined organic phases were washed with the indicated aqueous solution (if needed), dried over anhyd MgSO₄, filtered, and concentrated under reduced pressure using a rotary evaporator. The crude material was purified by flash chromatography using silica gel with the indicated eluent.

Methyl 8-Oxo-1,4-dioxaspiro[4.5]decane-7-carboxylate (2). A solution of **1** (3.0 g, 19.2 mmol) in Me₂CO₃ (20 mL) was added dropwise to a suspension of NaH (60% in oil, 1.54 g, 38.4 mmol) in Me₂CO₃ (10 mL) at rt. The reaction was heated to reflux for 4 h, then cooled to 0 °C. Saturated aq NH₄Cl (30 mL) was added, and the usual workup (EtOAc; brine) and purification (5–15% EtOAc in hexanes) afforded compound **2** (3.81 g, 93%) as a pale-yellow oil. Spectral data was consistent with that previously reported.⁴¹

Methyl 9-Hexyl-8-oxo-1,4-dioxaspiro[4.5]dec-6-ene-7-carboxylate (3). LDA (2.86 mmol) was prepared by the addition of *n*-BuLi (2.5 M in hexanes, 1.15 mL, 2.86 mmol) to a precooled (–78 °C) solution of diisopropylamine (402 μ L, 2.86 mmol) in THF (1.0 mL). The mixture was stirred for 30 min, then a solution of **2** (201.3 mg, 0.94 mmol) in THF (0.5 mL + 0.5 mL for rinsing) was added dropwise via cannula. The mixture was stirred for 1 h at 0 °C, then 1-iodohexane (0.69 mL, 4.68 mmol) was added. The resulting mixture was allowed to warm up to rt and stirred overnight. Water (15 mL) was added, and the usual workup (EtOAc) and purification (silica gel saturated with Et₃N, 0–5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (158.7 mg confirmed by mass spectrometry) as a yellow oil. A solution of the latter and pyridine (0.12 mL, 1.48 mmol) in CH₂Cl₂ (2.0 mL) was added to a solution of PhSeCl (269.9 mg, 1.41 mmol) in CH₂Cl₂ (2.0 mL) at rt. The mixture was stirred for 3 h, then 1 N aqueous HCl (10 mL) was added. The usual workup (CH₂Cl₂) and purification (10% Et₂O in hexanes) gave the corresponding phenylselenanyl- β -ketoester (197.3 mg, 46% over 2 steps) as a yellow oil. ¹H NMR (CDCl₃, 400 MHz) δ (ppm) 7.57 (d, 2H, *J* = 7.3 Hz), 7.40 (t, 1H, *J* = 7.3 Hz), 7.30 (t, 2H, *J* = 7.4 Hz), 3.89 (m, 4H), 3.65 (s, 3H), 2.93 (m, 1H), 2.50 (dd, 1H, *J* = 3.6, 13.7 Hz), 2.17 (d, 1H, *J* = 13.7 Hz), 2.01 (m, 1H), 1.88 (m, 1H), 1.79 (t, 1H, *J* = 13.5 Hz), 1.28 (br s, 9H), 0.87 (t, 3H, *J* = 6.7 Hz). ¹³C NMR (CDCl₃, 100 MHz) δ (ppm) 204.72, 170.07, 138.76, 129.68, 128.77, 126.38, 106.88, 64.93, 64.23, 59.69, 52.74, 46.61, 43.54, 41.75, 31.43, 29.41, 29.20, 26.80, 22.72, 14.22. HRMS (ESI-Q-ToF) calcd for C₂₂H₃₀O₅SeNa 477.1152, found 477.1152. IR (neat) ν 2952, 2926, 2856, 1713, 1436 cm^{–1}. An aqueous H₂O₂ solution (30%, 1.0 mL) was added to a solution of phenylselenanyl- β -ketoester (196.1 mg, 0.43 mmol) in CH₂Cl₂ (3.0 mL) at rt. The mixture was vigorously stirred for 90 min, then water (1 mL) and CH₂Cl₂ (3 mL) were added. The usual workup (CH₂Cl₂) afforded pure **3** (122.8 mg, 95%) as a pale-yellow oil. ¹H NMR (CDCl₃, 300 MHz) δ (ppm) 7.06 (d, 1H, *J* = 2.0 Hz), 4.07 (m, 4H), 3.81 (s, 3H), 2.72 (m, 1H), 2.21 (m, 1H), 2.05 (dd, 1H, *J* = 12.3, 13.2 Hz), 1.86 (m, 1H), 1.27 (br s, 9H), 0.87 (t, 3H, *J* = 6.8 Hz). ¹³C NMR (CDCl₃, 75 MHz) δ (ppm) 196.20, 164.59, 147.61, 132.55, 103.63, 65.31, 64.92, 52.31, 45.48, 38.37, 31.54, 29.14,

28.91, 26.39, 22.46, 13.96. HRMS (ESI-Q-ToF) calcd for $C_{16}H_{24}O_3Na$ + MeOH 351.1778, found 351.1785. IR (neat) ν 2926, 1746, 1436, 1256, 1083 cm^{-1} .

Methyl 9-Methyl-8-oxo-1,4-dioxaspiro[4.5]dec-6-ene-7-carboxylate (4). Following the procedure used to prepare 3, 2 (503.7 mg, 2.35 mmol) was treated with LDA (4.99 mmol) and iodomethane (0.16 mL, 2.57 mmol) then quenched with saturated aq NH_4Cl (15 mL) and water (2 mL). The usual workup (EtOAc) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (278.5 mg confirmed by mass spectrometry) as a pale-yellow oil. The latter was treated with pyridine (0.11 mL, 1.36 mmol) and PhSeBr (319.4 mg, 1.35 mmol) then quenched with 1 N aq HCl (10 mL). The usual workup (CH_2Cl_2 ; brine) and purification (20% EtOAc in hexanes) gave the corresponding phenylselenanyl- β -ketoester (109.2 mg, 32% over 2 steps) as a colorless oil. 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 7.58 (d, 2H, $J = 7.2$ Hz), 7.39 (t, 1H, $J = 7.3$ Hz), 7.30 (t, 2H, $J = 7.5$ Hz), 3.85 (m, 4H), 3.64 (s, 3H), 3.06 (m, 1H), 2.50 (dd, 1H, $J = 3.8, 13.7$ Hz), 2.15 (d, 1H, $J = 13.7$ Hz), 1.99 (m, 1H), 1.83 (t, 1H, $J = 13.4$ Hz), 1.13 (d, 3H, $J = 6.5$ Hz). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 170.17, 138.79, 129.71, 128.79, 126.46, 106.79, 64.97, 64.30, 59.43, 52.78, 43.69, 43.51, 40.75, 14.86. HRMS (ESI-Q-ToF) calcd for $C_{17}H_{20}O_5SeNa$ 407.0369, found 407.0371. IR (neat) ν 2896, 1736, 1709, 1219 cm^{-1} . Phenylselenanyl- β -ketoester (44.4 mg, 0.12 mmol) was treated with aq H_2O_2 (50 wt %, 0.1 mL), and then water (4 mL) and CH_2Cl_2 (10 mL) were added. The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) afforded pure 4 (27.1 mg, quant) as a pale-yellow oil. 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 7.07 (d, 1H, $J = 2.2$ Hz), 4.05 (m, 4H), 3.80 (s, 3H), 2.86 (m, 1H), 2.17 (ddd, 1H, $J = 2.2, 4.8, 13.5$ Hz), 2.07 (t, 1H, $J = 13.5$ Hz), 1.16 (d, 3H, $J = 6.7$ Hz). ^{13}C NMR ($CDCl_3$, 100 MHz) δ (ppm) 196.62, 164.77, 148.17, 132.62, 103.71, 65.52, 65.16, 52.58, 41.27, 40.83, 14.81. HRMS (ESI-Q-ToF) calcd for $C_{11}H_{14}O_5Na$ 249.0733, found 249.0740. IR (neat): 2955, 2934, 2891, 1744, 1690, 1435, 1251 cm^{-1} .

4,4-Dimethylcyclohexanone (5). A suspension of Pd/C (5 wt %, 729 mg, 0.4 mol %) in ethanol (20 mL) was added to a solution of 22⁴² (11.6 g, 93.4 mmol) in ethanol (60 mL) at rt. The mixture was cooled to 0 °C, the flask was purged with hydrogen, and the reaction was stirred at rt for 2 h. Complete conversion was assured by 1H NMR. The resulting mixture was filtered on Celite and washed with CH_2Cl_2 . The solvent was removed under vacuum to afford 5 (10.7 g, 91%) as a white solid. Spectral data was consistent with that previously reported.⁴³

Methyl 5,5-Dimethyl-2-oxocyclohexanecarboxylate (6). Following the procedure used to prepare 2, 5 (9.6 g, 76 mmol) was treated with NaH (60% in oil, 12.2 g, 305 mmol), Me_2CO_3 (16.0 mL, 190 mmol), and KH (30% in oil, catalytic amount) and then quenched with aq 3 M AcOH (120 mL) and poured into brine (100 mL). The usual workup (EtOAc, 5 $\times$ 100 mL) followed by distillation (<1 Torr, 74–76 °C) afforded 6 (10.1 g, 72%) as a colorless oil. Spectral data was consistent with that previously reported.⁴⁴

Methyl 3,3-Dimethyl-6-oxocyclohex-1-enecarboxylate (7). Following the procedure used to prepare 3, 6 (102.6 mg, 0.56 mmol) was treated with pyridine (60 μ L, 0.74 mmol) and PhSeBr (163.5 mg, 0.69 mmol). The usual workup (CH_2Cl_2) gave the corresponding phenylselenanyl- β -ketoester. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then water (10 mL) was added. The usual workup (CH_2Cl_2 ; aqueous saturated $NaHCO_3$; brine) and purification (10–20% EtOAc in hexanes) afforded 7 (58.2 mg, 57%) as a yellow oil. Spectral data was consistent with that previously reported.⁴³

Methyl 5-Hexyl-3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (8). Following the procedure used to prepare 3, 6 (204.1 mg, 1.11 mmol) was treated with LDA (2.40 mmol) in THF (1.0 mL) and 1-iodohexane (0.25 mL, 1.69 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (256.8 mg confirmed by mass spectrometry) as a yellow oil. A solution of the latter (49.7 mg, 0.19 mmol) in THF (0.3 mL + 0.2 mL for rinsing) was added to a solution of DDQ (57.2 mg, 0.25 mmol) in THF (0.5 mL) via canula at rt. The mixture was stirred for 24 h, and

then water (2 mL) was added. The usual workup (EtOAc; brine) and purification (10% EtOAc in hexanes) afforded 8 (19.8 mg, 34% over two steps) as a yellow oil. 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 7.25 (d, 1H, $J = 2.0$ Hz), 3.79 (s, 3H), 2.46 (m, 1H), 1.90 (m, 2H), 1.65 (t, 1H, $J = 13.7$ Hz), 1.29 (m, 9H), 1.24 (s, 3H), 1.20 (s, 3H), 0.87 (t, 3H, $J = 6.7$ Hz). ^{13}C NMR ($CDCl_3$, 100 MHz) δ (ppm) 197.09, 165.79, 163.13, 130.24, 52.35, 43.59, 41.83, 33.91, 31.87, 30.36, 29.53, 29.03, 26.88, 25.40, 22.77, 14.24. HRMS (ESI-Q-ToF) calcd for $C_{16}H_{26}O_3Na$ 289.1774, found 289.1784. IR (neat) ν 2955, 2925, 2857, 1745, 1719, 1686 cm^{-1} .

Methyl 3,3,5-Trimethyl-6-oxocyclohex-1-enecarboxylate (9). Following the procedure used to prepare 3, 6 (215.8 mg, 1.17 mmol) was treated with LDA (2.34 mmol) and iodomethane (80 μ L, 1.29 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (150.0 mg, confirmed by mass spectrometry) as a yellow oil. The latter (100.0 mg, 0.50 mmol) was treated with pyridine (50 μ L, 0.62 mmol) and PhSeBr (140.1 mg, 0.59 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) gave the corresponding phenylselenanyl- β -ketoester as a yellow oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol). The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) and purification (5–10% EtOAc in hexanes) afforded 9 (58.4 mg, 39% over two steps) as a yellow oil. 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 7.25 (s, 1H), 3.78 (s, 3H), 2.62 (m, 1H), 1.83 (ddd, 1H, $J = 2.3, 4.9, 13.5$ Hz), 1.69 (t, 1H), 1.25 (s, 3H), 1.18 (s, 3H), 1.12 (d, 3H, $J = 6.6$ Hz). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 197.31, 165.72, 163.38, 129.98, 52.30, 44.61, 38.67, 33.97, 30.28, 25.43, 14.87. HRMS (ESI-Q-ToF) calcd for $C_{11}H_{16}O_3$ 219.992, found 219.1000. IR (neat) ν 2962, 1743, 1718, 1686, 1285, 1258 cm^{-1} .

Methyl 5-Allyl-3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (10). Following the procedure used to prepare 3, 6 (203.0 mg, 1.10 mmol) was treated with LDA (2.40 mmol) and allyl bromide (0.15 mL, 1.73 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc) and purification (10% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (210.7 mg, confirmed by mass spectrometry) as a yellow oil. Following the procedure used to prepare 8, alkylated β -ketoester (101.5 mg, 0.45 mmol) was treated with DDQ (57.2 mg, 0.25 mmol) and K_2CO_3 (76.5 mg, 0.55 mmol) and then quenched with water (4 mL). The usual workup (EtOAc; brine) and purification (5–10% EtOAc in hexanes) afforded 10 (45.9 mg, 39% over two steps) as a yellow oil. Spectral data was consistent with that previously reported.⁴⁵

Methyl 5-Benzyl-3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (11). Following the procedure used to prepare 3, 6 (203.9 mg, 1.11 mmol) was treated with LDA (2.40 mmol) and benzyl bromide (0.20 mL, 1.68 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (303.4 mg, confirmed by mass spectrometry) as a yellow oil. Following the procedure used to prepare 8, alkylated β -ketoester (102.6 mg, 0.37 mmol) was treated with DDQ (97.8 mg, 0.43 mmol) in 1,4-dioxane, and then hexanes were added, the suspension was filtered, and the solvent was evaporated. The usual and purification (5–10% EtOAc in hexanes) afforded 11 (40.5 mg, 40% over two steps) as a pale-brown oil. 1H NMR (CD_2Cl_2 , 400 MHz) δ (ppm) 7.22 (m, 6H), 3.73 (s, 3H), 3.35 (dd, 1H, $J = 4.0, 14.0$ Hz), 2.78 (m, 1H), 2.43 (dd, 1H, $J = 9.0, 14.0$ Hz), 1.71 (m, 1H), 1.59 (t, 1H, $J = 13.7$ Hz), 1.12 (s, 3H), 1.11 (s, 3H). ^{13}C NMR (CD_2Cl_2 , 100 MHz) δ (ppm) 196.17, 165.32, 163.32, 140.22, 130.23, 129.46, 128.86, 126.37, 52.27, 45.45, 41.36, 35.35, 33.99, 30.17, 25.02. HRMS (ESI-Q-ToF) calcd for $C_{17}H_{20}O_3$ 295.1305, found 295.1314. IR (neat) ν 2958, 1742, 1717, 1690, 1271, 1258 cm^{-1} .

Methyl 3,3-Dimethyl-6-oxo-5-(prop-2-yn-1-yl)cyclohex-1-enecarboxylate (12). Following the procedure used to prepare 3, 6 (199.3 mg, 1.08 mmol) was treated with LDA (2.41 mmol) and propargyl bromide (80% in toluene, 185 μ L, 1.66 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc) and purification (5% EtOAc in hexanes) gave the corresponding alkylated

β -ketoester (176.8 mg confirmed by mass spectrometry) as a yellow oil. Following the procedure used to prepare **8**, alkylated β -ketoester (98.0 mg, 0.44 mmol) was treated with DDQ (159.7 mg, 0.70 mmol) and K_2CO_3 (165.7 mg, 1.20 mmol) in 1,4-dioxane and then quenched with water (10 mL). The usual workup (EtOAc; brine) and purification (10% EtOAc in hexanes) afforded **12** (44.7 mg, 34% over two steps) as a white solid. 1H NMR ($CDCl_3$, 400 MHz) δ (ppm) 7.34 (d, 1H, J = 2.2 Hz), 3.80 (s, 3H), 2.81 (ddd, 1H, J = 2.7, 4.0, 17.0 Hz), 2.72 (m, 1H), 2.30 (ddd, 1H, J = 2.6, 8.7, 17.0 Hz), 2.17 (ddd, 1H, J = 2.3, 4.6, 13.4 Hz), 1.98 (t, 1H, J = 2.7 Hz), 1.78 (t, 1H, J = 13.8 Hz), 1.29 (s, 3H), 1.24 (s, 3H). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 194.59, 165.30, 164.37, 129.58, 81.98, 70.11, 52.43, 42.82, 41.09, 34.03, 30.27, 25.31, 18.86. HRMS (ESI-Q-ToF) calcd for $C_{13}H_{16}O_3$ 243.0992, found 243.0995. IR (neat) ν 3262, 2967, 1716, 1684, 1259 cm^{-1} .

Methyl 3,3-Dimethyl-6-oxo-5-(2,2,2-trifluoroethyl)cyclohex-1-enecarboxylate (13). Following the procedure used to prepare **3**, **6** (203.8 mg, 1.11 mmol) was treated with LDA (2.42 mmol) and 1,1,1-trifluoro-2-iodoethane (130 μ L, 1.33 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (136.3 mg) as a yellow oil. The latter (136.3 mg) was treated with pyridine (50 μ L, 0.62 mmol) and PhSeBr (138.1 mg, 0.59 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) afforded the corresponding phenylselanyl- β -ketoester as an orange oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol). The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) and purification (5% EtOAc in hexanes) afforded **13** (20.1 mg, 7% over two steps) as a yellow oil. 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 7.35 (d, 1H, J = 2.3 Hz), 3.81 (s, 3H), 3.09 (m, 1H), 2.88 (m, 1H), 2.14 (ddd, 1H, J = 13.0, 4.2, 1.7 Hz), 1.94 (m, 1H), 1.76 (t, 1H, J = 13.9 Hz), 1.30 (s, 3H), 1.23 (s, 3H). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 193.23, 164.99, 164.10, 129.22, 125.40, 52.51, 41.93, 38.76 (q, J = 2.2 Hz), 33.97, 32.74 (q, J = 28.9 Hz), 30.18, 24.99. HRMS (ESI-Q-ToF) calcd for $C_{12}H_{15}F_3O_3$ 287.0866, found 287.0870. IR (neat) ν 2957, 1745, 1717, 1690, 1252 cm^{-1} .

Methyl 5-(2-(tert-Butoxy)-2-oxoethyl)-3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (14). Following the procedure used to prepare **3**, **6** (210.4 mg, 1.14 mmol) was treated with LDA (2.47 mmol) and *tert*-butyl bromoacetate (250 μ L, 1.69 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (5–20% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (164.0 mg confirmed by mass spectrometry) as a yellow oil. The latter (93.5 mg) was treated with pyridine (61 μ L, 0.75 mmol) and PhSeBr (175.4 mg, 0.74 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) afforded the corresponding phenylselanyl- β -ketoester as a yellow oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then water (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) and purification (10–20% EtOAc in hexanes) afforded **14** (36.0 mg, 19% over two steps) as a white solid; mp = 47–49 °C. 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 7.30 (d, 1H, J = 2.2 Hz), 3.79 (s, 3H), 3.10–2.97 (m, 1H), 2.86 (dd, 1H, J = 16.6, 5.2 Hz), 2.15 (dd, 1H, J = 16.6, 7.4 Hz), 1.90 (ddd, 1H, J = 13.3, 4.8, 2.2 Hz), 1.76 (t, 1H, J = 13.6 Hz), 1.46 (s, 9H), 1.29 (s, 3H), 1.21 (s, 3H). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 195.07, 171.62, 165.37, 163.92, 129.56, 80.82, 52.41, 41.93, 40.84, 35.52, 34.08, 30.26, 28.25, 25.16. HRMS (ESI-Q-ToF) calcd for $C_{16}H_{24}O_5Na$ 319.1516, found 319.1525. IR (neat) ν 2976, 1724, 1364, 1147, 1125 cm^{-1} .

Methyl 5-(Cyanomethyl)-3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (15). Following the procedure used to prepare **3**, **6** (828.8 mg, 4.50 mmol) was treated with LDA (9.88 mmol) and bromoacetonitrile (0.63 mL, 9.04 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (5–20% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (145.2 mg confirmed by mass spectrometry) as an orange oil. The latter (139.0 mg) was treated with pyridine (0.29 mL, 3.13 mmol) and PhSeBr (588.4 mg, 2.49 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine)

afforded the corresponding phenylselanyl- β -ketoester as a brown oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then water (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) and purification (20–30% EtOAc in hexanes) afforded **15** (84.8 mg, 9% over two steps) as a yellow oil. 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 7.40 (d, 1H, J = 2.2 Hz), 3.80 (s, 3H), 2.91 (m, 2H), 2.49 (dd, 1H, J = 18.1, 9.5 Hz), 2.15 (ddd, 1H, J = 13.4, 4.4, 2.2 Hz), 1.86 (t, 1H, J = 13.6 Hz), 1.33 (s, 3H), 1.26 (s, 3H). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 192.61, 164.93, 164.62, 128.92, 118.22, 52.56, 41.21, 40.67, 34.19, 30.12, 25.25, 17.96. HRMS (ESI-Q-ToF) calcd for $C_{12}H_{13}NO_3Na$ 244.0944, found 244.0950. IR (neat) ν 2959, 2249, 1737, 1684, 1275, 1256, 1218 cm^{-1} .

Methyl 3,3-Dimethyl-6-oxo-5-(2-oxopropyl)cyclohex-1-enecarboxylate (16). Following the procedure used to prepare **3**, **6** (801.1 mg, 4.35 mmol) was treated with LDA (9.62 mmol) and chloroacetone (0.69 mL, 8.67 mmol) and then quenched with aq 1 N HCl (30 mL). The usual workup (EtOAc; brine) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (215.4 mg confirmed by mass spectrometry) as an orange oil. The latter (171.7 mg) was treated with pyridine (0.33 mL, 3.56 mmol) and PhSeBr (680.1 mg, 2.88 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) afforded the corresponding phenylselanyl- β -ketoester as a brown oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then water (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) and purification (20–30% EtOAc in hexanes) afforded **16** (37.8 mg, 4% over two steps) as a white solid; mp = 53–55 °C. 1H NMR ($CDCl_3$, 300 MHz) δ (ppm) 7.29 (d, 1H, J = 2.3 Hz), 3.80 (s, 3H), 2.69 (dd, 1H, J = 14.6, 4.5 Hz), 2.64 (d, 1H, 4.4 Hz), 2.55 (d, 1H, J = 4.4 Hz), 1.89 (ddd, 1H, J = 13.5, 4.5, 2.3 Hz), 1.57 (t, 1H, 14.0 Hz), 1.47 (s, 3H), 1.25 (s, 3H), 1.22 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz) δ (ppm) 194.21, 165.23, 163.69, 130.12, 55.98, 52.41, 50.94, 49.20, 37.46, 33.92, 30.33, 25.04, 21.87. HRMS (ESI-Q-ToF) calcd for $C_{13}H_{18}O_4Na$ 261.1097, found 261.1113. IR (neat) ν 2944, 1709, 1682, 1265, 1226 cm^{-1} .

Methyl 5-(3-Methoxy-3-oxopropyl)-3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (17). Following the procedure used to prepare **3**, **6** (201.3 mg, 1.09 mmol) was treated with LDA (3.50 mmol) and methyl acrylate (0.15 mL, 1.67 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (2–20% EtOAc in hexanes) afforded the corresponding alkylated β -ketoester (162.6 mg confirmed by mass spectrometry) as an orange oil. The latter (143.3 mg) was treated with pyridine (0.17 mL, 1.84 mmol) and PhSeBr (400.1 mg, 1.70 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) afforded the corresponding phenylselanyl- β -ketoester as a brown oil. The latter was treated with aqueous H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then water (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq $NaHCO_3$; brine) and purification (5–30% EtOAc in hexanes) afforded **17** (40.7 mg, 16% over two steps) as a colorless oil. 1H NMR (CD_2Cl_2 , 300 MHz) δ (ppm) 7.20 (s, 1H), 3.72 (s, 3H), 3.62 (s, 3H), 2.55 (m, 1H), 2.39 (m, 2H), 2.12 (m, 1H), 1.84 (m, 1H), 1.68 (d, 1H, J = 13.8 Hz), 1.56 (m, 1H), 1.22 (s, 3H), 1.17 (s, 3H). ^{13}C NMR ($CDCl_3$, 75 MHz) δ (ppm) 196.29, 173.71, 165.19, 162.79, 130.00, 51.97, 51.46, 42.63, 41.88, 33.82, 31.45, 29.94, 24.91, 24.70. HRMS (ESI-Q-ToF) calcd for $C_{14}H_{20}O_5Na$ 291.1203, found 291.1211. IR (neat) ν 2954, 1733, 1685, 1267, 1218, 1168 cm^{-1} .

Allyl 3,3-Dimethyl-6-oxocyclohex-1-enecarboxylate (18). In a sealed tube equipped with a septum, DMAP (213.3 mg, 1.75 mmol) and 4 Å molecular sieves (500.8 mg) were added to a solution of **6** (219.2 mg, 1.19 mmol) in allyl alcohol (5.0 mL). The tube was sealed, the mixture was refluxed for 48 h, allowed to cool down to rt, and then aqueous 1 N HCl (10 mL) was added. The usual workup (EtOAc; brine) and purification (2% EtOAc in hexanes) gave a mixture of **6** (0.09 mmol) and the corresponding allyl ester (0.68 mmol) as a pale-yellow oil (158.1 mg). The mixture was dissolved in 1,4-dioxane (2.0 mL), and then K_2CO_3 (169.9 mg, 1.23 mmol) and DDQ (233.1 mg, 1.02 mmol) were added. The reaction mixture was heated to 100 °C for 4 h, cooled to rt, and quenched with water (10 mL). The usual workup (EtOAc; brine) and purification (5–10% EtOAc in hexanes)

afforded **18** (78.8 mg, 32% over two steps) as an orange oil. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.34 (s, 1H), 5.96 (m, 1H), 5.38 (ddd, 1H, $J = 1.5, 3.0, 17.2$ Hz), 5.26 (ddd, 1H, $J = 1.3, 2.6, 10.4$ Hz), 4.70 (dt, 2H, $J = 1.4, 5.7$ Hz), 2.53 (t, 2H, $J = 6.8$ Hz), 1.89 (t, 2H, $J = 6.78$ Hz), 1.23 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 194.40, 164.58, 164.36, 131.99, 130.24, 118.76, 65.88, 35.59, 35.35, 33.64, 27.51. HRMS (ESI-Q-ToF) calcd for $\text{C}_{12}\text{H}_{16}\text{O}_3\text{Na}$ 231.992, found 231.1003. IR (neat) ν 2962, 1740, 1714, 1686, 1267, 1224 cm^{-1} .

tert-Butyl 3,3-Dimethyl-6-oxocyclohex-1-enecarboxylate (19). Following the procedure used to prepare **18**, **6** (204.7 mg, 1.11 mmol) was treated DMAP (200.7 mg, 1.64 mmol) and 4 Å molecular sieves (365.8 mg) in *tert*-butanol (5.0 mL) and then was quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (silica gel saturated with Et_3N , 100% hexanes) gave a mixture of **6** (0.36 mmol) and the corresponding *tert*-butyl ester (0.40 mmol) as a colorless oil (156.1 mg). Following the procedure used to prepare **3**, the mixture was treated with pyridine (155 μL , 1.67 mmol) and PhSeBr (375.4 mg, 1.59 mmol) and then quenched with 1 N aq HCl (10 mL). The usual workup (CH_2Cl_2 ; 1 N aq HCl; brine) afforded the corresponding phenylselenanyl- β -ketoester as a brown oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then aq saturated NaHCO_3 (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq NaHCO_3) and purification (10–20% EtOAc in hexanes) afforded **19** (61.7 mg, 25% over two steps) as a white solid; mp = 28–30 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.17 (s, 1H), 2.50 (t, 2H, $J = 6.8$ Hz), 1.86 (t, 2H, $J = 6.8$ Hz), 1.50 (s, 9H), 1.21 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 194.73, 164.23, 162.53, 131.73, 81.83, 35.69, 35.41, 33.43, 28.23, 27.59. HRMS (ESI-Q-ToF) calcd for $\text{C}_{13}\text{H}_{20}\text{O}_3\text{Na}$ 247.1305, found 247.1311. IR (neat) ν 2957, 1725, 1673, 1367, 1290, 1237 cm^{-1} .

3,3-Dimethyl-6-oxocyclohex-1-enecarboxylic Acid (20). A solution of KOH (85%, 894.0 mg, 15.9 mmol) in water (1.5 mL) was added to a solution of **7** (54.0 mg, 0.30 mmol) in MeOH (5.0 mL) at rt. The mixture was refluxed for 1 h, cooled to rt, diluted with EtOAc (30 mL), and quenched with aqueous 3 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (30% EtOAc in hexanes) afforded **20** (24.8 mg, 49%) as a white solid; mp = 68–70 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 12.47 (s, 1H), 8.11 (s, 1H), 2.66 (m, 2H), 1.94 (m, 2H), 1.27 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 202.86, 173.76, 164.08, 124.53, 35.05, 34.61, 34.35, 27.01. HRMS (ESI-Q-ToF) calcd for $\text{C}_9\text{H}_{12}\text{O}_3\text{Na}$ 191.0679, found 191.0677. IR (neat) ν 2960, 2761, 1742, 1637, 1607, 1439, 1223, 1156 cm^{-1} .

Methyl 6-Hydroxy-3,3-dimethylcyclohex-1-enecarboxylate (21). CeCl_3 (heptahydrate, 189.3 mg, 0.51 mmol) and then NaBH_4 (20.6 mg, 0.54 mmol) were added to a solution of **7** (91.8 mg, 0.50 mmol) in MeOH (2.0 mL) at rt. The solution was stirred (gas evolution) for 20 h, and then aq 1 N HCl (2 mL) was added dropwise followed by water (3 mL). The usual workup (EtOAc) and purification (10–20% EtOAc in hexanes) afforded **21** (53.2 mg) as a pale-yellow oil. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 6.76 (s, 1H), 4.48 (t, 1H, $J = 5.1$ Hz), 3.77 (s, 3H), 3.71 (s, 1H), 1.93–1.75 (m, 2H), 1.64 (ddd, 1H, $J = 13.5, 9.9, 3.6$ Hz), 1.42 (m, 1H), 1.11 (s, 3H), 1.03 (s, 3H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 176.21, 168.22, 151.90, 129.89, 71.59, 63.87, 51.96, 51.92, 47.48, 40.75, 37.24, 33.22, 32.25, 32.21, 30.46, 29.72, 28.85, 27.74, 27.25, 24.31. HRMS (ESI-Q-ToF) calcd for $\text{C}_{10}\text{H}_{16}\text{O}_3\text{Na}$ 207.0992, found 207.0993. IR (neat) ν 3445, 2949, 2865, 1702, 1361, 1281, 1251 cm^{-1} .

2-Chloro-4,4-dimethylcyclohex-2-enone (23). (Diacetoxyiodo)benzene (320.0 mg, 0.99 mmol) and pyridine hydrochloride (238.9 mg, 2.07 mmol) were added to a solution of **22** (100.1 mg, 0.81 mmol) in CH_2Cl_2 (4.0 mL) at rt. The mixture was stirred for 6 h, and then (diacetoxyiodo)benzene (334.0 mg, 1.04 mmol) and pyridine hydrochloride (233.6 mg, 2.02 mmol) were added again. The mixture was stirred overnight and then quenched with aq 1 N HCl (10 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl) and purification (5% EtOAc in hexanes) afforded **23** (32.7 mg, 25%) as a yellow oil. Spectral data was consistent with that previously reported.⁴⁶

2-Iodo-4,4-dimethylcyclohex-2-enone (24). A solution of iodine (868.1 mg, 3.42 mmol) and pyridine (3.0 mL) in CCl_4 (3.0

mL) was added dropwise to a solution of **22** (233.7 mg, 1.88 mmol) and pyridine (3.0 mL) in CCl_4 (3.0 mL) at 0 °C. The resulting mixture was allowed to warm up to rt, stirred for 3 h, and then quenched with aq 1 N HCl (10 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; aq saturated NaHCO_3) and purification (5% EtOAc in hexanes) afforded **24** (351.6 mg, 75%) as a yellow oil. Spectral data was consistent with that previously reported.⁴⁷

2-(Hydroxymethyl)-4,4-dimethylcyclohex-2-enone (25). A solution of **22** (1.6 g, 12.9 mmol) in THF (2.0 mL + 2.0 mL for rinsing) was added to a mixture of paraformaldehyde (194.6 mg, 6.4 mmol) and imidazole (441.6 mg, 6.5 mmol) in 1 N NaHCO_3 (25.6 mL) and THF (2.4 mL) at rt. The mixture was stirred for 50 h and then quenched with aq 1 N HCl (25 mL). The usual workup (CH_2Cl_2) and purification (10–30% EtOAc in hexanes) afforded 4,4-dimethylcyclohex-2-enone (548.2 mg, 4.4 mmol) and **25** (453.8 mg, 35% corrected) as a colorless oil. Spectral data was consistent with that previously reported.⁴⁸

(3,3-Dimethyl-6-oxocyclohex-1-en-1-yl)methyl 5,5-dimethyl-2-oxocyclohexanecarboxylate (26). A solution of **25** (129.5 mg, 0.84 mmol) in toluene (2.0 mL + 2 $\times$ 1.0 mL for rinsing) was added dropwise to a solution of **6** (300.4 mg, 1.63 mg) in toluene (5.0 mL) at rt. Triethylamine (0.23 mL, 1.65 mmol) was then added, and the resulting mixture was refluxed in a Dean–Stark apparatus for 24 h. The solution was allowed to cool down to rt, washed with brine (3 $\times$ 5 mL), dried over anhyd Na_2SO_4 , filtered, and concentrated. The usual purification (2–5% EtOAc in hexanes) afforded methyl 5,5-dimethyl-2-oxocyclohexanecarboxylate (120.8 mg) and **26** (241.1 mg, 94% corrected) as a colorless oil. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 12.12 (s, 1H), 6.61 (s, 1H), 4.78 (d, 2H, $J = 0.9$ Hz), 2.50 (m, 2H), 2.29 (t, 2H, $J = 6.7$ Hz), 2.03 (s, 2H), 1.87 (t, 2H, $J = 6.8$ Hz), 1.44 (t, 2H, $J = 6.7$ Hz), 1.18 (s, 6H), 0.95 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 198.08, 197.86, 172.44, 171.72, 169.96, 157.95, 156.99, 131.57, 131.07, 96.49, 61.90, 61.01, 53.82, 42.06, 39.40, 38.08, 36.13, 36.00, 34.60, 34.49, 33.15, 33.10, 30.92, 30.22, 29.11, 27.99, 27.92, 27.85, 27.81, 26.79, 24.73. HRMS (ESI-Q-ToF) calcd for $\text{C}_{18}\text{H}_{26}\text{O}_4\text{Na}$ 329.1723, found 329.1732. IR (neat) ν 2969, 1669, 1634, 1606, 1230, 1200, 1174 cm^{-1} .

Methyl 5,5-Dimethyl-2-oxocyclohex-3-enecarboxylate (27). Following the procedure used to prepare **2**, **22** (534.0 mg, 4.30 mmol) was treated with NaH (60% in oil, 505.5 mg, 12.6 mmol) and Me_2CO_3 (1.7 mL, 20.2 mmol) in refluxing 1,4-dioxane (3 mL) overnight and then quenched with water (2 mL) and aq 3 M AcOH (2 mL). The usual workup (Et_2O) and purification (5% EtOAc in hexanes) gave **27** (265.9 mg, 34%) as a yellow oil. The spectral data was consistent with that previously reported.⁴⁹

Methyl 1,5,5-Trimethyl-2-oxocyclohex-3-enecarboxylate (28). Iodomethane (30 μL , 0.48 mmol) was added dropwise to a solution of **27** (54.5 mg, 0.30 mmol) in acetone (3.0 mL) at 0 °C, followed by K_2CO_3 (115.0 mg, 0.83 mmol). The resulting mixture was stirred overnight at 40 °C, was allowed to cool down to rt, and the solvent was removed under vacuum. Aq 1 N HCl (5 mL) was added, and then the usual workup (CH_2Cl_2 ; brine) and purification (5% EtOAc in hexanes) afforded **28** (31.7 mg, 54%) as a yellow oil. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 6.59 (dd, 1H, $J = 1.9, 10.2$ Hz), 5.95 (d, 1H, $J = 10.2$ Hz), 3.68 (s, 3H), 2.43 (dd, 1H, $J = 1.9, 14.2$ Hz), 1.81 (d, 1H, $J = 14.2$ Hz), 1.38 (s, 3H), 1.14 (s, 3H), 1.07 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz) δ (ppm) 197.16, 174.89, 157.96, 126.31, 52.61, 51.24, 47.37, 33.40, 31.47, 26.84, 23.07. HRMS (ESI-Q-ToF) calcd for $\text{C}_{11}\text{H}_{16}\text{O}_3\text{Na}$ 219.992, found 219.993. IR (neat) ν 2958, 1731, 1682, 1249, 1228 cm^{-1} .

Methyl 3,3-Dimethyl-6-oxocyclohexa-1,4-dienecarboxylate (29). Selenium dioxide (193.8 mg, 1.75 mmol) was added to a solution of **27** (104.0 mg, 0.57 mmol) in *tert*-butanol (5 wt % AcOH, 10.0 mL) at 30 °C. The resulting mixture was stirred at reflux for 20 h and allowed to cool down to rt. Et_2O (10 mL) was added, layers were separated, and the organic layer was washed with aq saturated NaHCO_3 (3 $\times$ 5 mL). The usual workup (Et_2O ; brine) and purification (10–50% EtOAc in hexanes) gave the product contaminated with selenium. This material was vaporized under vacuum with a heat gun, and the vapor was collected in another flask

to afford pure **29** (50.0 mg, 49%) as a yellow oil. Spectral data was consistent with that previously reported.⁴⁹

Dimethyl 5,5'-(Methylenebis(3,3-dimethyl-6-oxocyclohex-1-enecarboxylate)) (30). Following the procedure used to prepare **3, 6** (400.5 mg, 2.17 mmol) was treated with LDA (4.80 mmol) and diiodomethane (105 μ L, 1.30 mmol) and then quenched with saturated aq 1 N HCl (20 mL). The usual workup (EtOAc; brine) and purification (10% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (374.8 mg confirmed by mass spectrometry) as an orange oil. The latter (102.3 mg, 0.27 mmol) was treated with pyridine (70 μ L, 0.87 mmol) and PhSeBr (192.9 mg, 0.82 mmol) and then quenched with aq 1 N HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) afforded the corresponding phenylselanyl- β -ketoester as an orange oil. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol), and then saturated aq NaHCO_3 (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq NaHCO_3 ; brine) and purification (20–25% EtOAc in hexanes) afforded **30** (28.9 mg, 25% over two steps) as a yellowish solid; mp = 132–134 °C. ^1H NMR (CD_2Cl_2 , 400 MHz) δ (ppm) 7.19 (br s, 2H), 3.71 (s, 6H), 2.79 (m, 2H), 1.88 (m, 2H), 1.66 (m, 4H), 1.26 (s, 6H), 1.17 (s, 6H). ^{13}C NMR (CD_2Cl_2 , 100 MHz) δ (ppm) 197.69, 165.48, 163.03, 130.25, 52.22, 43.18, 42.13, 34.20, 30.68, 30.18, 25.27. HRMS (ESI-Q-ToF) calcd for $\text{C}_{21}\text{H}_{28}\text{O}_6\text{Na}$ 399.1778, found 399.1785. IR (neat) ν 2967, 1709, 1692, 1259, 1235 cm^{-1} .

Dimethyl 5,5'-(Propane-1,3-diyl)bis(3,3-dimethyl-6-oxocyclohex-1-enecarboxylate) (31). Following the procedure used to prepare **3, 6** (404.1 mg, 2.19 mmol) was treated with LDA (4.80 mmol) and 1,3-diiodopropane (150 μ L, 1.31 mmol) and then quenched with saturated aq NH_4Cl (20 mL). The usual workup (EtOAc; brine) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (355.8 mg confirmed by mass spectrometry) as a yellow oil. The latter (100.8 mg, 0.25 mmol) was treated with pyridine (90 μ L, 1.11 mmol) and PhSeBr (237.2 mg, 1.01 mmol) and then quenched with 1 N aq HCl (5 mL). The usual workup (CH_2Cl_2 ; brine) gave the corresponding phenylselanyl- β -ketoester. The latter was treated with aq H_2O_2 (50 wt %, 0.5 mL, 8.65 mmol). The usual workup (CH_2Cl_2 ; saturated aq NaHCO_3 ; brine) and purification (20–25% EtOAc in hexanes) afforded **31** (63.0 mg, 50% over two steps) as a yellow oil. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.27 (s, 2H), 3.79 (s, 6H), 2.50 (m, 2H), 1.89 (ddd, 4H, J = 2.2, 4.7, 13.4 Hz), 1.67 (t, 2H, J = 18.3 Hz), 1.39 (m, 4H), 1.25 (s, 6H), 1.21 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 197.01, 196.97, 165.67, 163.30, 163.27, 130.16, 52.33, 52.30, 43.46, 43.43, 42.02, 41.97, 33.98, 30.33, 29.44, 29.30, 25.38, 24.30, 24.23. HRMS (ESI-Q-ToF) calcd for $\text{C}_{23}\text{H}_{32}\text{O}_6\text{Na}$ 427.2091, found 427.2109. IR (neat) ν 2955, 1742, 1717, 1682, 1270, 1215 cm^{-1} .

5,5'-(Butane-1,4-diyl)bis(3,3-dimethyl-6-oxocyclohex-1-enecarboxylate) (32). Following the procedure used to prepare **3, 6** (201.3 mg, 1.09 mmol) was treated with LDA (2.50 mmol) and 1,4-diiodobutane (87 μ L, 0.66 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (199.5 mg confirmed by mass spectrometry) as a white gum. The latter (106.7 mg, 0.25 mmol) was treated with pyridine (90 μ L, 1.11 mmol) and PhSeBr (233.8 mg, 0.99 mmol) and then quenched with aq 1 N HCl (5 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl; brine) gave the corresponding phenylselanyl- β -ketoester as a yellow oil. The latter was treated with aq H_2O_2 (50 wt %, 1.0 mL, 17.3 mmol), and then saturated aq NaHCO_3 (5 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq NaHCO_3 ; brine) and purification (30% EtOAc in hexanes) afforded **32** (64.3 mg, 53% over two steps) as a white solid; mp = 84–86 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.25 (d, 2H, J = 2.1 Hz), 3.79 (s, 6H), 2.48 (m, 2H), 1.89 (m, 4H), 1.65 (td, 2H, J = 0.9, 13.9 Hz), 1.34 (m, 6H), 1.25 (s, 6H), 1.20 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 196.97, 165.74, 163.19, 130.22, 52.34, 43.58, 41.90, 33.92, 30.35, 29.05, 29.03, 27.17, 27.14, 25.40. HRMS (ESI-Q-ToF) calcd for $\text{C}_{24}\text{H}_{34}\text{O}_6\text{Na}$ 441.2248, found 441.2251. IR (neat) ν 2936, 1717, 1675, 1259, 1237 cm^{-1} .

(E)-Dimethyl 5,5'-(But-2-ene-1,4-diyl)bis(3,3-dimethyl-6-oxocyclohex-1-enecarboxylate) (33). Following the procedure used

to prepare **3, 6** (201.3 mg, 1.09 mmol) was treated with LDA (2.50 mmol) and (2E)-1,4-dibromobut-2-ene (146.7 mg, 0.69 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (10% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (201.9 mg confirmed by mass spectrometry) as a yellow oil. Following the procedure used to prepare **8**, alkylated β -ketoester (89.9 mg, 0.21 mmol) was treated with DDQ (106.5 mg, 0.47 mmol) in 1,4-dioxane and then quenched with saturated aq Na_2CO_3 (10 mL). The mixture was poured into brine (10 mL). The usual workup (EtOAc) and purification (10–30% EtOAc in hexanes) gave a yellow solid that was triturated in Et_2O to obtain **33** (12.9 mg, 13% over two steps) as a white solid; mp = 126–128 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.27 (d, 2H, J = 2.2 Hz), 5.43 (m, 2H), 3.79 (s, 6H), 2.54 (m, 4H), 2.08 (m, 2H), 1.85 (m, 2H), 1.61 (td, 2H, J = 3.7, 13.6 Hz), 1.23 (s, 6H), 1.19 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 196.19, 196.16, 165.61, 163.55, 130.08, 130.05, 129.95, 52.35, 43.53, 43.51, 41.39, 41.37, 33.96, 33.95, 32.33, 32.31, 30.33, 25.34. HRMS (ESI-Q-ToF) calcd for $\text{C}_{24}\text{H}_{32}\text{O}_6\text{Na}$ 439.2091, found 439.2094. IR (neat) ν 2963, 1717, 1674, 1273, 1252 cm^{-1} .

Dimethyl 5,5'-(But-2-yne-1,4-diyl)bis(3,3-dimethyl-6-oxocyclohex-1-enecarboxylate) (34). Following the procedure used to prepare **3, 6** (214.7 mg, 1.17 mmol) was treated with LDA (2.64 mmol), freshly distilled HMPA (200 μ L, 1.15 mmol), and 1,4-dichloro-2-butyne (64 μ L, 0.65 mmol) and then quenched with aq 1 N HCl (10 mL). The usual workup (EtOAc; brine) and purification (5% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (127.4 mg confirmed by mass spectrometry) as a yellow oil. Following the procedure used to prepare **8**, alkylated β -ketoester (121.4 mg, 0.29 mmol) was treated with DDQ (150.1 mg, 0.66 mmol) in 1,4-dioxane and then quenched with saturated aq Na_2CO_3 (10 mL). The usual workup (EtOAc; brine) and purification (20–40% EtOAc in hexanes) gave a brown residue that was triturated in Et_2O to afford **34** (17.5 mg, 8% over two steps) as a light-brown solid; mp = 132–134 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.31 (d, 2H, J = 2.1 Hz), 3.78 (s, 6H), 2.68 (m, 4H), 2.27 (m, 2H), 2.11 (m, 2H), 1.75 (t, 2H, J = 13.7 Hz), 1.26 (s, 6H), 1.23 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 194.89, 194.87, 165.36, 165.34, 164.22, 164.15, 129.71, 129.67, 79.25, 79.24, 52.35, 43.19, 43.17, 41.21, 41.19, 33.97, 30.27, 25.33, 19.16. HRMS (ESI-Q-ToF) calcd for $\text{C}_{24}\text{H}_{30}\text{O}_6\text{Na}$ 437.1935, found 437.1938. IR (neat) ν 2956, 1731, 1670, 1281, 1255 cm^{-1} .

Bis(methyl 8-oxo-1,4-dioxaspiro[4.5]dec-6-ene-7-carboxylate) Methylene (35). Following the procedure used to prepare **3, 2** (196.7 mg, 0.92 mmol) was treated with LDA (1.93 mmol) and dibromomethane (32 μ L, 0.46 mmol) and then quenched with aq 1 N HCl (10 mL) and water (2 mL). The usual workup (Et_2O ; brine) and purification (30% EtOAc in hexanes) gave the corresponding alkylated β -ketoester (52.6 mg confirmed by mass spectrometry) as a pale-yellow solid; mp = 148–150 °C. The latter (47.5 mg, 0.11 mmol) was treated with pyridine (50 μ L, 0.62 mmol) and PhSeBr (107.6 mg, 0.46 mmol) and then quenched with aq 1 N HCl (3 mL). The usual workup (CH_2Cl_2 ; aq 1 N HCl) and purification (20–40% EtOAc in hexanes) gave the corresponding phenylselanyl- β -ketoester (37.7 mg, 13% over 2 steps) as a white solid. The latter (23.1 mg, 0.03 mmol) was treated with aq H_2O_2 (50 wt %, 0.05 mL), and then water (1 mL) was added. The usual workup (CH_2Cl_2 ; saturated aq NaHCO_3 ; brine) afforded pure **36** (15.0 mg, quant) as a brown solid; mp = 196 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.07 (d, 2H, J = 1.8 Hz), 4.06 (m, 8H), 3.80 (s, 6H), 3.04 (m, 2H), 2.44 (m, 2H), 2.12 (d, 1H, J = 29.2 Hz), 2.07 (d, 1H, J = 25.8 Hz), 1.81 (t, 2H, J = 6.4 Hz). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 196.72, 164.62, 148.48, 132.53, 103.69, 65.64, 65.23, 52.62, 44.22, 39.81, 30.39. HRMS (ESI-Q-ToF) calcd for $\text{C}_{21}\text{H}_{24}\text{O}_{10}\text{Na}$ 459.1262, found 459.1269. IR (neat) ν 2956, 1742, 1717, 1690, 1257, 1219 cm^{-1} .

Bis(methyl 8-oxo-1,4-dioxaspiro[4.5]dec-6-ene-7-carboxylate) Propylene (36). Following the procedure used to prepare **3**, a solution of **2** (101.7 mg, 0.48 mmol) in THF (1.0 mL) was treated with a previously prepared solution of LDA (1.0 mmol) in THF (1.0 mL) and then with 1,3-diiodopropane (34 μ L, 0.29 mmol). Aqueous 1 N HCl (3 mL) was added, and the usual workup (EtOAc; brine) and purification (10–20% EtOAc in hexanes) gave the corresponding

alkylated β -ketoester (63.7 mg confirmed by mass spectrometry) as a colorless oil. A solution of PhSeCl (106.9 mg, 0.56 mmol) in CH_2Cl_2 (1.0 mL) was treated with pyridine (39 μL , 0.48 mmol) at rt for 30 min, and then with a solution of the alkylated β -ketoester (54.5 mg, 0.12) in CH_2Cl_2 (1.0 mL) at rt for 8 h. CH_2Cl_2 (4 mL), water (1 mL), and 1 N aqueous HCl (1 mL) were added, and layers were separated. The usual workup (CH_2Cl_2 ; brine) and purification (0–20% EtOAc in hexanes) afforded the corresponding phenylselenanyl- β -ketoester (62.7 mg confirmed by mass spectrometry, 38% over 2 steps) as a pale-yellow solid. A solution of the latter (60.5 mg, 0.08 mmol) in CH_2Cl_2 (1.0 mL) was treated with aqueous H_2O_2 (30 wt %, 0.04 mL). Water (5 mL) was added, and the usual workup (CH_2Cl_2 ; brine) afforded 37 (37.1 mg, quant) as a white solid; mp = 138–140 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.06 (d, 2H, J = 1.9 Hz), 4.18–3.95 (m, 8H), 3.81 (s, 6H), 2.83–2.66 (m, 1H), 2.22 (ddd, 2H, J = 13.5, 4.7, 2.0 Hz), 2.13–1.99 (m, 2H), 1.97–1.77 (m, 2H), 1.49–1.32 (m, 4H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 196.21, 164.79, 148.05, 132.80, 103.88, 65.60, 65.20, 52.59, 45.60, 38.67, 29.22, 23.84. HRMS (ESI-Q-ToF) calcd for $\text{C}_{23}\text{H}_{28}\text{O}_{10}\text{Na}$ 487.1575, found 487.1582. IR (neat) ν 2952, 2891, 1741, 1683, 1436, 1260 cm^{-1} .

(3,3-Dimethyl-6-oxocyclohex-1-en-1-yl)methyl 3,3-dimethyl-6-oxocyclohex-1-enecarboxylate (37). K_2CO_3 (110.0 mg, 0.80 mmol) and DDQ (103.5 mg, 0.46 mmol) were added to a solution of 26 (97.2 mg, 0.32 mmol) in 1,4-dioxane (2.0 mL). The solution was stirred at reflux for 4 h and allowed to cool down to rt. Water (10 mL) was added, and then the usual workup (EtOAc; brine) and purification (10–20% EtOAc in hexanes) afforded 37 (72.9 mg, 75%) as a white solid; mp = 40–42 °C. ^1H NMR (CDCl_3 , 300 MHz) δ (ppm) 7.32 (app s, 1H), 6.84 (app s, 1H), 4.85 (d, 2H, J = 1.3 Hz), 2.51 (m, 4H), 1.88 (m, 4H), 1.22 (s, 6H), 1.19 (s, 6H). ^{13}C NMR (CDCl_3 , 75 MHz) δ (ppm) 198.07, 194.52, 164.79, 164.57, 157.47, 131.45, 130.30, 62.12, 36.04, 35.59, 35.36, 34.64, 33.67, 33.15, 27.90, 27.50. HRMS (ESI-Q-ToF) calcd for $\text{C}_{18}\text{H}_{24}\text{O}_4\text{Na}$ 327.1567, found 327.1575. IR (neat) ν 2956, 1710, 1667, 1269, 1227, 1177, 1160 cm^{-1} .

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.jmedchem.6b01132.

NQO1 induction data, LDH assay data, NMR spectra for new compounds (PDF)

Molecular formula strings (CSV)

■ AUTHOR INFORMATION

Corresponding Author

*Phone: 819-821-7822. Fax: 819-821-8017. E-mail: Guillaume.Belanger@USherbrooke.ca.

Author Contributions

L.J.D. synthesized new compounds, H.T. tested for biological activity, L.J.D., G.B., and E.M. designed compounds, H.T. and M.R. designed biological experiments, and all authors reviewed and approved manuscript contents. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The Institut de Pharmacologie de Sherbrooke is acknowledged for a scholarship to LD. M.V.R. is a FRQS Junior 2 Research Scholar. Funding by Centre de Recherche du Centre Hospitalier de l'Université de Sherbrooke, Faculté des sciences de l'Université de Sherbrooke.

■ ABBREVIATIONS USED

Nrf2, nuclear factor erythroid 2-related factor 2; Keap1, Kelch-like ECH-associated protein 1; ARE, antioxidant response element; HEK, human embryonic kidney; ROS, reactive oxygen species; RNS, reactive nitrogen species; FDA, food and drug administration; tBHQ, *tert*-butyl hydroquinone; CDDO, 2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oic acid; BTB, broad complex, tramtrack, bric-a-brac; IVR, intervening region; LDA, lithium diisopropylamine; DDQ, 2,3-dichloro-5,6-dicyano-*p*-benzoquinone; LDH, lactate dehydrogenase; SEM, standard error of the mean; GSH, glutathione; clogP, calculated partition coefficient

■ REFERENCES

- (1) Lichtenberg, D.; Pinchuk, I. Oxidative Stress, the Term and the Concept. *Biochem. Biophys. Res. Commun.* **2015**, *461*, 441–444.
- (2) Suzuki, T.; Motohashi, H.; Yamamoto, M. Toward Clinical Application of the Keap1-Nrf2 Pathway. *Trends Pharmacol. Sci.* **2013**, *34*, 340–346.
- (3) O'Connell, M. A.; Hayes, J. D. The Keap1/Nrf2 Pathway in Health and Disease: From the Bench to the Clinic. *Biochem. Soc. Trans.* **2015**, *43*, 687–689.
- (4) Osburn, W. O.; Kensler, T. W. Nrf2 Signaling: An Adaptive Response Pathway for Protection against Environmental Toxic Insults. *Mutat. Res., Rev. Mutat. Res.* **2008**, *659*, 31–39.
- (5) Hayes, J. D.; Dinkova-Kostova, A. T. The Nrf2 Regulatory Network Provides an Interface between Redox and Intermediary Metabolism. *Trends Biochem. Sci.* **2014**, *39*, 199–218.
- (6) Hancock, R.; Schaap, M.; Pfister, H.; Wells, G. Peptide Inhibitors of the Keap1-Nrf2 Protein-Protein Interaction with Improved Binding and Cellular Activity. *Org. Biomol. Chem.* **2013**, *11*, 3553–3557.
- (7) Marcotte, D.; Zeng, W.; Hus, J. C.; McKenzie, A.; Hession, C.; Jin, P.; Bergeron, C.; Lugovskoy, A.; Enyedy, I.; Cuervo, H.; et al. Small Molecules Inhibit the Interaction of Nrf2 and the Keap1 Kelch Domain through a Non-Covalent Mechanism. *Bioorg. Med. Chem.* **2013**, *21*, 4011–4019.
- (8) Jnoff, E.; Albrecht, C.; Barker, J. J.; Barker, O.; Beaumont, E.; Bromidge, S.; Brookfield, F.; Brooks, M.; Bubert, C.; Ceska, T.; et al. Binding Mode and Structure-Activity Relationships around Direct Inhibitors of the Nrf2-Keap1 Complex. *ChemMedChem* **2014**, *9*, 699–705.
- (9) Satoh, M.; Saburi, H.; Tanaka, T.; Matsuura, Y.; Naitow, H.; Shimozono, R.; Yamamoto, N.; Inoue, H.; Nakamura, N.; Yoshizawa, Y.; Aoki, T.; Tanimura, R.; Kunishima, N. Multiple Binding Modes of a Small Molecule to Human Keap1 Revealed by X-Ray Crystallography and Molecular Dynamics Simulation. *FEBS Open Bio* **2015**, *5*, 557–570.
- (10) Zhuang, C.; Narayanapillai, S.; Zhang, W.; Sham, Y. Y.; Xing, C. Rapid Identification of Keap1-Nrf2 Small-Molecule Inhibitors through Structure-Based Virtual Screening and Hit-Based Substructure Search. *J. Med. Chem.* **2014**, *57*, 1121–1126.
- (11) Zheng, S.; Santosh Laxmi, Y. R.; David, E.; Dinkova-Kostova, A. T.; Shrivani, K. H.; Ren, Y.; Zheng, Y.; Trevino, I.; Bumeister, R.; Ojima, I.; Wigley, W. C.; Bliska, J. B.; Mierke, D. F.; Honda, T. Synthesis, Chemical Reactivity as Michael Acceptors, and Biological Potency of Monocyclic Cyanoenones, Novel and Highly Potent Anti-Inflammatory and Cytoprotective Agents. *J. Med. Chem.* **2012**, *55*, 4837–4846.
- (12) Dinkova-Kostova, A. T.; Talalay, P.; Sharkey, J.; Zhang, Y.; Holtzclaw, W. D.; Wang, X. J.; David, E.; Schiavoni, K. H.; Finlayson, S.; Mierke, D. F.; Honda, T. An Exceptionally Potent Inducer of Cytoprotective Enzymes: Elucidation of the Structural Features That Determine Inducer Potency and Reactivity with Keap1. *J. Biol. Chem.* **2010**, *285*, 33747–33755.
- (13) Honda, T.; Yoshizawa, H.; Sundarajan, C.; David, E.; Lajoie, M. J.; Favalaro, F. G.; Janosik, T.; Su, X.; Honda, Y.; Roebuck, B. D.; Gribble, G. W. Tricyclic Compounds Containing Nonenolizable

Cyano Enones. A Novel Class of Highly Potent Anti-Inflammatory and Cytoprotective Agents. *J. Med. Chem.* **2011**, *54*, 1762–1778.

(14) Satoh, T.; McKercher, S. R.; Lipton, S. A. Reprint of: Nrf2/ARE-Mediated Antioxidant Actions of pro-Electrophilic Drugs. *Free Radical Biol. Med.* **2014**, *66*, 45–57.

(15) Xi, M. Y.; Sun, Z. Y.; Sun, H. P.; Jia, J. M.; Jiang, Z. Y.; Tao, L.; Ye, M.; Yang, X.; Wang, Y. J.; Xue, X.; Huang, J.; Gao, Y.; Guo, X.; Zhang, S.; Yang, Y.; Guo, Q.; Hu, R.; You, Q. Synthesis and Bioevaluation of a Series of α -Pyrone Derivatives as Potent Activators of Nrf2/ARE Pathway (Part I). *Eur. J. Med. Chem.* **2013**, *66*, 364–371.

(16) Wu, K. C.; McDonald, P. R.; Liu, J.; Klaassen, C. D. Screening of Natural Compounds as Activators of the Keap1-Nrf2 Pathway. *Planta Med.* **2014**, *80*, 97–104.

(17) Venci, J. V.; Gandhi, M. A. Dimethyl Fumarate (Tecfidera): A New Oral Agent for Multiple Sclerosis. *Ann. Pharmacother.* **2013**, *47*, 1697–1702.

(18) Li, W.; Zheng, S.; Higgins, M.; Morra, R. P., Jr.; Mendis, A. T.; Chien, C.-W.; Ojima, I.; Mierke, D. F.; Dinkova-Kostova, A. T.; Honda, T. New Monocyclic, Bicyclic, and Tricyclic Ethynylcyanodienones as Activators of the Keap1/Nrf2/ARE Pathway and Inhibitors of Inducible Nitric Oxide Synthase. *J. Med. Chem.* **2015**, *58*, 4738–4748.

(19) Baird, L.; Llères, D.; Swift, S.; Dinkova-Kostova, A. T. Regulatory Flexibility in the Nrf2-Mediated Stress Response Is Conferred by Conformational Cycling of the Keap1-Nrf2 Protein Complex. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110*, 15259–15264.

(20) Cleasby, A.; Yon, J.; Day, P. J.; Richardson, C.; Tickle, I. J.; Williams, P. A.; Callahan, J. F.; Carr, R.; Concha, N.; Kerns, J. K.; Qi, H.; Sweitzer, T.; Ward, P.; Davies, T. G. Structure of the BTB Domain of Keap1 and Its Interaction with the Triterpenoid Antagonist CDDO. *PLoS One* **2014**, *9*, e98896.

(21) Kobayashi, M.; Li, L.; Iwamoto, N.; Nakajima-Takagi, Y.; Kaneko, H.; Nakayama, Y.; Eguchi, M.; Wada, Y.; Kumagai, Y.; Yamamoto, M. The Antioxidant Defense System Keap1-Nrf2 Comprises a Multiple Sensing Mechanism for Responding to a Wide Range of Chemical Compounds. *Mol. Cell. Biol.* **2009**, *29*, 493–502.

(22) Flanagan, M. E.; Abramite, J. A.; Anderson, D. P.; Aulabaugh, A.; Dahal, U. P.; Gilbert, A. M.; Li, C.; Montgomery, J.; Oppenheimer, S. R.; Ryder, T.; et al. Chemical and Computational Methods for the Characterization of Covalent Reactive Groups for the Prospective Design of Irreversible Inhibitors. *J. Med. Chem.* **2014**, *57*, 10072–10079.

(23) Bauer, R. A. Covalent Inhibitors in Drug Discovery: From Accidental Discoveries to Avoided Liabilities and Designed Therapies. *Drug Discovery Today* **2015**, *20*, 1061–1073.

(24) Several reports stress the importance of having two electrophilic sites on a molecule to improve its activity toward the Keap1/Nrf2/ARE pathway. However, none of these reports systematically evaluated the length and rigidity of the spacer between these electrophiles, or the effect of the electrophilicity of the acceptors. See ref 12. See also: (a) Dinkova-Kostova, A. T.; Massiah, M. A.; Bozak, R. E.; Hicks, R. J.; Talalay, P. Potency of Michael Reaction Acceptors as Inducers of Enzymes That Protect against Carcinogenesis Depends on Their Reactivity with Sulfhydryl Groups. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, *98*, 3404–3409. (b) Turpaev, K.; Ermolenko, M.; Cresteil, T.; Drapier, J. C. Benzylidenemalononitrile Compounds as Activators of Cell Resistance to Oxidative Stress and Modulators of Multiple Signaling Pathways. A Structure–activity Relationship Study. *Biochem. Pharmacol.* **2011**, *82*, 535–547.

(25) Barker, D.; Brimble, M. A.; McLeod, M. D.; Savage, G. P. Synthesis of Tricyclic Analogues of Methyllycaconitine Using Ring Closing Metathesis to Append a B Ring to an AE Azabicyclic Fragment. *Org. Biomol. Chem.* **2004**, *2*, 1659–1669.

(26) Liotta, D.; Barnum, C.; Puleo, R.; Zima, G.; Bayer, C.; Kezar, H. S., III A Simple Method for the Efficient Synthesis of Unsaturated β -Dicarbonyl Compounds. *J. Org. Chem.* **1981**, *46*, 2920–2923.

(27) With a 1,1-dicarbonyl, these Michael acceptors are often referred to as Nazarov acceptors.

(28) It should be noted that, due to the instability of final compounds **3** and **4**, purification had to be performed after selenation. Oxidation furnished clean products.

(29) Boddaert, T.; Coquerel, Y.; Rodriguez, J. Combination of Rearrangement with Metallic and Organic Catalyses – a Step- and Atom-Economical Approach to α -Spiroactones and -Lactams. *Eur. J. Org. Chem.* **2011**, *2011*, 5061–5070.

(30) Magee, D. I.; Ratshonka, S.; Mcconaghy, J.; Hood, M. Synthesis of β - and β,β -Substituted Morita–Baylis–Hillman Adducts Using a Two-Step Protocol. *Can. J. Chem.* **2012**, *90*, 450–463.

(31) Ngatimin, M.; Gartshore, C. J.; Kindler, J. P.; Naidu, S.; Lupton, D. W. The α -Halogenation of α,β -Unsaturated Carbonyls and Dihalogenation of Alkenes Using Bisacetoxiodobenzene/pyridine Hydrohalides. *Tetrahedron Lett.* **2009**, *50*, 6008–6011.

(32) Pandey, G.; Balakrishnan, M. Suzuki Cross-Coupling/Reductive Debenzyloxycarbonylation Sequence for the Syntheses of [c] Annulated Isoquinolines: Application for the Syntheses of Pancratisatin-like Isoquinolines. *J. Org. Chem.* **2008**, *73*, 8128–8131.

(33) Williams, D. R.; Reeves, J. T.; Nag, P. P.; Pitcock, W. H.; Baik, M. H. Studies of the Generation and Pericyclic Behavior of Cyclic Pentadienyl Carbanions. Alkylation Reactions as an Efficient Route to Functionalized *cis*-Bicyclo[3.3.0]octenes. *J. Am. Chem. Soc.* **2006**, *128*, 12339–12348.

(34) Mhasni, O.; Rezgui, F. The First Et_3N -Mediated Transesterification of β -Keto Esters Using Baylis–Hillman Alcohols. *Tetrahedron* **2011**, *67*, 6322–6326.

(35) Anxionnat, B.; Robert, B.; George, P.; Ricci, G.; Perrin, M.-A.; Gomez Pardo, D.; Cossy, J. Ring Expansion of Cyclic β -Amino Alcohols Induced by Diethylaminosulfur Trifluoride: Synthesis of Cyclic Amines with a Tertiary Fluorine at C3. *J. Org. Chem.* **2012**, *77*, 6087–6099.

(36) Liu, H.-J.; Browne, E. N. C. Total Synthesis of α - and β -Himachalene by an Intermolecular Diels–Alder Approach. *Can. J. Chem.* **1981**, *59*, 601–608.

(37) As suggested by Satoh et al., hydrophilic electrophiles are more readily to react with the cysteine of glutathione (GSH) because of its hydrophilic microenvironment. Depletion of GSH could explain the activity we observed: on one hand, Nrf2 transcription might be activated to restore normal GSH concentration. On the other hand, the compound would also be more inclined to react with Keap1 owing to depletion of GSH pool. See ref 14.

(38) In fact, Honda reported **29** with $\text{CO}_2\text{Me}=\text{CN}$ as a very potent modulator of Keap1/Nrf2. See ref 11.

(39) Dinkova-Kostova, A. T.; Holtzclaw, W. D.; Cole, R. N.; Itoh, K.; Wakabayashi, N.; Katoh, Y.; Yamamoto, M.; Talalay, P. Direct Evidence That Sulfhydryl Groups of Keap1 Are the Sensors Regulating Induction of Phase 2 Enzymes That Protect against Carcinogens and Oxidants. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, *99*, 11908–11913.

(40) Baell, J. B.; Holloway, G. A. New Substructure Filters for Removal of Pan Assay Interference Compounds (PAINS) from Screening Libraries and for Their Exclusion in Bioassays. *J. Med. Chem.* **2010**, *53*, 2719–2740.

(41) Lavallée, J.-F.; Spino, C.; Ruel, R.; Hogan, K. T.; Deslongchamps, P. *Can. J. Chem.* **1992**, *70*, 1406–1426.

(42) Romanski, S.; Kraus, B.; Guttentag, M.; Schlundt, W.; Rücker, H.; Adler, A.; Neudörfl, J.-M.; Alberto, R.; Amslinger, S.; Schmalz, H.-G. *Dalt. Trans.* **2012**, *41*, 13862–13875.

(43) Liu, H.-J.; Browne, E. N. C.; Chew, S. Y. *Can. J. Chem.* **1988**, *66*, 2345–2347.

(44) Madder, A.; De Clercq, P. J.; Declercq, J.-P. *J. Org. Chem.* **1998**, *63*, 2548–2559.

(45) Chou, H. H.; Wu, H. M.; Wu, J. D.; Ly, T. W.; Jan, N. W.; Shia, K. S.; Liu, H. J. *Org. Lett.* **2008**, *10*, 121–123.

(46) Altmeyer, I.; Margaretha, P. *Helv. Chim. Acta* **1977**, *60*, 874–881.

(47) Soorukram, D.; Knochel, P. *Org. Lett.* **2004**, *6*, 2409–2411.

(48) Williams, D. R.; Reeves, J. T.; Nag, P. P.; Pitcock, W. H.; Baik, M. H. *J. Am. Chem. Soc.* **2006**, *128*, 12339–12348.

(49) Liu, H.-J.; Browne, E. N. C. *Can. J. Chem.* **1981**, *59*, 601–608.